

## STUDIES ON MICRO PROPAGATION OF BANANA, CV. BARGIBALE

HEGDE, A. K.,<sup>1,2</sup> HEGDE, V. K.<sup>1,2</sup> AND HEGDE, V.<sup>2</sup>

<sup>1</sup>MES's M. M. Arts and Science College, Sirsi, Uttara Kannada, Karnataka; <sup>2</sup>Bhandimane Life-Science Research Foundation, Nejjur Colony, Banavasi road, Sirsi, Uttara Kannada (Karnataka).  
E. mail: [vinayhegde.bhandimane@gmail.com](mailto:vinayhegde.bhandimane@gmail.com), Cell: 09538652458

Received: February 17, 2018; Accepted: March 11, 2018

**Abstract:** *In the present study, an efficient reproducible protocol for high frequency micropropagation of Banana Var. Bargibale was described. An in-vitro protocol for Cv. Bargibale was established by using shoot tip culture. The healthy suckers which were collected from the disease free and high yielding mother plants were surfaced sterilized and initiated on standard MS medium supplemented with different combinations and concentrations of plant growth hormones. Out of tried combinations, maximum multiplication (90%) was obtained in MS medium supplemented with 3.5 mg/L BAP, 10 mg/L adenine sulphate and 0.5 mg/L IAA, while maximum elongation of shoot (3.6cm) was observed in MS medium supplemented with 3 mg/L BAP, 8 mg/L adenine sulphate and 0.5 mg/L IAA. Then the micro shoots were transferred to in-vitro rooting medium MS medium half strength, supplemented with 1 mg/L BAP, 2.5 mg/L IAA and 2 g/L activated charcoal. Then excellent acclimatization was observed for transferred plantlets with 90% survival rate.*

**Key words:** Banana Cv. Bargibale, Micro propagation

### INTRODUCTION

Banana is one of the most important fruit crop in the world, both as a staple food as well as a major export commodity for many tropical and sub-tropical countries [1,2]. Uttara Kannada has rich biodiversity supporting number of ecological functions as well as livelihood of thousands of peoples in the district. Banana is one of the chiefly available fruits in the Uttara Kannada district. There are number of local varieties of banana cultivated in the region viz., Boodibale, Kembale, Karibale, Metlibale Mysore Metli, Nenibale, Baragibale, Rasabale, Pachebale, Sakkarebale and Shanbale etc. To safeguard sustainable banana production and generate wealth for small-scale farmers, high quality planting material is crucial. The extensive basic work on the *in vitro* propagation of banana [3,4] had led to the technological development of *in vitro* mass production

of different cultivars.

Plant tissue culture techniques can potentially overcome some of the factors limiting traditional approaches to banana improvement. Banana is generally propagated vegetatively through suckers. But the traditional method is laborious; time consuming and not very efficient as far as production of homogenous plant is concerned [5]. Only 5 to 10 suckers can be obtained from a plant per year in conventional method. Furthermore, banana production sometimes becomes seriously affected by different diseases [6]. As a result, banana productivity decreases and the yield become very poor. To overcome this problem, production of saplings using *in vitro* culture techniques could be an effective for production of planting materials of bananas. A large number of uniform disease free plants can be produced from a single plant or even a small plant

tissue (explants) showing good genetic potential in this method [7] and plant multiplication can be continued throughout the year irrespective of seasonal variation. Tissue cultured plants grow vigorously, establish more quickly and take a shorter time to bunch emergence and harvest.

Tissue culture technique produce 39% higher yield than conventional sword suckers [8]. The application of tissue culture and rapid propagation method for banana production continues to become more widely used in both developed and developing countries. Tissue culture techniques can be applied not only to increase propagation rates but also to modify the germ plasm itself. Banana is successfully micropropagated through plant tissue culture [9,10].

## MATERIALS AND METHODS

**Plant materials and surface sterilization of explants:** Fresh suckers of Bargibale were collected from farmer's fields of Uttara Kannada district with identification. The suckers were chopped off about 3-5 cm length and washed thoroughly under running tap water for 10-15 min. Shoot tips were prepared by trimming corm and outer leaf sheaths from the suckers. These shoot tips were collected in a beaker containing tap water, then brought to the laboratory and washed thoroughly with tap water, then washed with detergent mixed with Tween 80 and finally with distilled water for three times. These shoot tips were treated with 0.1%  $\text{HgCl}_2$  solution for different periods (4,6,8 and 10 min) to determine time period for contamination free culture. They were then washed with sterile distilled water for 5 minutes, three times. Treatment of shoot tips with 0.1%  $\text{HgCl}_2$  for 10 min was found suitable for sterilization and finally washed thoroughly with sterile distilled water under aseptic conditions in a laminar air flow cabinet.

**Culture conditions and hardening:** All explants were placed on autoclaved Murashige and Skoog [MS medium, 1962] medium with different concentrations of cytokinin (1, 1.5, 2, 2.5, 3, 3.5, 4, 4.5 and 5mg/l BAP), auxin (0.5 mg/l IAA) and adenine sulphate (0, 2, 4, 6, 8, 10, 12, 14 and 16 mg/l). The pH of medium was adjusted to 5.8 prior to autoclaving. 3% sucrose was supplemented in the media. The culture bottles containing explants were incubated in a temperature controlled growth chamber. 10 bottle, replication based on explants were incubated at  $25\pm 2^\circ\text{C}$  with 12-16 hr photoperiod

(approximately 2000 lux) provided by white fluorescent tubes. The materials were sub cultured at a regular interval of four weeks into same medium to produce multiple shoots. For *in-vitro* rooting, elongated shoots were transferred to a rooting medium, half strength Murashige and Skoog medium with different concentrations of auxin (1, 1.5, 2, 2.5, 3, 3.5, 4mg/l IAA), cytokinin BAP 1 mg/L and 2 g/L activated charcoal for four weeks maintaining standard culture conditions. After four to five weeks, the plantlets were carefully removed from the culture vessels. The roots of the plantlets were gently washed under running tap water to remove agar attached to the roots. The plantlets were hardened under poly tunnel for 4 weeks and shade house for 6 weeks.

## RESULTS

The results of the study showed that shoot tips obtained from the sword suckers of banana cv. Bargibale were cultured onto MS medium supplemented with (1, 1.5, 2, 2.5, 3, 3.5, 4, 4.5 and 5mg/l BAP), (0.5 mg/l IAA) and adenine sulphate (0, 2, 4, 6, 8, 10, 12, 14 and 16 mg/l) for induction of multiple shoot (Figs. a-d). After two weeks of culture for organogenesis, all culture was refreshed on same media. After four weeks of sub culture, the percentage of shoot proliferation under various concentrations of BAP was found in the range of 65 to 90%. Out of tried concentrations, MS medium supplemented with BAP at a concentration of 3.5 mg/L +0.5 mg/L IAA and Adenine Sulphate 10 mg/L yielded best response on shoot proliferation which accounted to 90%. BAP at a concentration of 1.0 mg/l yielded lowest response on shoot proliferation, which was 65%. During the culture period, the number of shoots increased gradually in all the media (Table 1) and (Figs. a-d). MS medium supplemented with BAP at a concentration of 3 mg/L + 0.5 mg/L IAA and adenine sulphate 8 mg/L yielded best shoot elongation, which was found to be 3.6 cm in shoot length. While MS medium supplemented with BAP at a concentration of 5 mg/l+0.5 mg/l IAA and adenine sulphate 16 mg/l yielded the lowest shoot elongation, which was found to be 2.6 cm in shoot length (Table 1).

The regenerated shoots were cultured on half strength of MS medium supplemented with different concentrations of auxin (1, 1.5, 2, 2.5, 3mg/l IAA), 1mg/l BAP and 2g/l activated charcoal for *in-vitro* rooting. Although few roots developed from multiple



**Fig: 1:** a) Bargibale (selected banana cultivar) suckers were aseptically cultured on MS medium. b and c) Sub culturing for Multiplication and Shoot elongation. d) In-vitro rooting

shoots during culture, but those roots are not so healthy for future growth and development of plantlets. IAA at a concentration of 2.5 mg/l was found to be the best medium for *in-vitro* rooting in banana cv. Bargibale (Table 2) and (Fig. d). Well-rooted plantlets were hardened under poly tunnel for 4 weeks and shade house for 6 weeks. The polythene bags were covered with a plastic film to retain moisture and gradually acclimatized to outdoor condition. The final survival rate of the transferred plantlets to soil was 90% and their growth in such condition was satisfactory.

## DISCUSSION

Tissue culture technique involves the establishment of different cell or tissue under a suitable culture

condition, *in vitro* cell proliferation and subsequent regeneration of plants [11]. In banana, the most widespread used technique for propagation is reproduction by taking actively growing piece of stems called as micro cuttings for *in vitro* micro propagation under varying concentration of different cytokinins and auxins [12,13]. The present investigation was carried out for the induction and proliferation of multiple shoot from shoot tip and subsequent rooting. Different concentrations of cytokinin and auxin were supplied with MS medium for shoot and root induction. However, some report showed that MS medium with 0.5 mg/l BAP +0.05 mg/l NAA +10% 9(v/v) CW was optimum for maximum number of shoot regenerated from sucker explants [14] longest shoot in banana when cultured on MS medium added with 25  $\mu$ M BAP was obtained

**Table 1:** Effect of different concentrations of BAP and Adenine sulphate on shoot multiplication of Banana Cv. Bargibale.

Concentration in mg/L			Response %	Number of shoots/explants	Shoot length (in cm)
BAP	Adenine sulphate	IAA			
1.0	0	0.5	65	3	3.6
1.5	2	0.5	75	3	3.5
2.0	4	0.5	80	5	3.5
2.5	6	0.5	80	5	3.6
3.0	8	0.5	90	7	3.6
3.5	10	0.5	90	8	3.4
4.0	12	0.5	80	8	3.1
4.5	14	0.5	75	7	2.8
5.0	16	0.5	60	5	2.6

[15]. MS medium [16] in concentration with 4 mg/l BAP + 2.0 mg/l NAA + 13% CW was optimum for highest number of shoot regeneration in banana from sucker explants whereas others [17] developed a medium for regeneration of banana from excised floral apices which was MS +2.0 mg/l BAP+ 1.0 mg/l kin +1.0 mg/l IAA +15% CW.

In our observation, maximum multiplication (90%) was obtained in MS medium supplemented with 3.5 mg/L BAP, 10 mg/L adenine sulphate and 0.5 mg/L IAA, while maximum elongation of shoot (3.6cm) was observed in MS medium supplemented with 3 mg/L BAP, 8 mg/L adenine sulphate and 0.5 mg/L IAA. The best *in-vitro* rooting was found in MS medium half strength, supplemented with 1 mg/L BAP, 2.5 mg/L IAA and 2 g/L activated charcoal. Then excellent acclimatization was observed for transferred plantlets with 90% survival rate.

### ACKNOWLEDGEMENTS

The authors are grateful to UGC SWRO for financial support through Minor research project. The authors are grateful to the staff members of Department of Biotechnology, the Principal of MES's M.M. Arts and Science College Sirsi. The authors are grateful to the members and founders of Bhandimane Life Science Research Foundation, Sirsi, Karnataka, India.

### REFERENCES

- [1] Singh, H.P.: *Indian bananas-Issues and strategies. Global conference on Banana and Plantain*, October 28-31, Bangalore, India. Abstracts, 1-2 (2002).
- [2] Bose, T.K.: Banana. In: *Fruits of India- Tropical and sub-tropical*, Naya Prakash Publication, Calcutta, India, pp 124-161 (1985).

- [3] Kodym, A. and Zapata, F.A.: *Plant Cell Tissue Org. Cult.* 55: 141-145 (1999).
- [4] Nandwani, D., Zehr, U., Zehr, B.E. and Barwale, R.B.: *Gartenbauwissenschaft.* 65: 237-240 (2000).
- [5] Banaerjee, N. and, De Langhe, E.: *Plant Cell Rep.* 4: 351-354 (1985).
- [6] Rahman, M.Z., Nasiruddin, K.M., Amin, M.A. and Islam, M.N.: *Asian J. Plant Sci.*; 3: 406-409 (2004).
- [7] Martin, K.P., Pachathundikandi, S.K., Zhang, C.L., Slater, A. and Joseph, M.: *In Vitro Cell. Dev. Biol. Plant.* 42: 188-192 (2006).
- [8] Farahani, F., Aminpoor, H., Sheidai, M.Z., Noormahammadi, Z. and Mazinani, M.H.: *Asian J. Plant Sci.* 7: 116-118 (2008).
- [9] Bhatt, P. and Bhatt, D.: *Banana Micropropagation- A success story. Comprehensive Micropropagation of Horticultural Crops.* International Book Distributing Co., Lucknow, pp 212-220 (2003).
- [10] Vuylsteke, D. and DE Langhe, E.A.L.: *Trop. Agri. (Trinidad)*, 62: 323-328 (1985).
- [11] Vasil, I.K., Scowcroft, W.R. and Frey, K.J.: *Plant Improvement and Somatic Cell Genetics*, Acad. Press, NY. 158-178 (1982).
- [12] Mendes, B.J.M., Filippi, S.B., Demetrio, G.B. and Rodriguez, A.P.M.: *Plant Cell Rep.* 18: 967-971 (1999).
- [13] Vuylsteke, D.R. and Ortiz, R.: *Hort. Sci.* 31: 862-865 (1996).
- [14] Akbar, M.A., Hossain, M.Z. and Roy, S.K.: *Bangladesh J. Life Sci.*, 15: 33-38 (2003).
- [15] Khanam, D., Hoque, M.A., Khan, M.A. and Quasem, A.: *Plant Tissue Cult.* 6: 89-94 (1996).
- [16] Habiba, U. and Reza, S.: *Plant Tissue Cult.* 12: 117-124 (2002).
- [17] Azad, M.A.K. and Amin, M.N.: *Plant Tissue Cult.* 11: 1-9 (2001).